

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number
WO 03/014080 A2

(51) International Patent Classification⁷: **C07D 207/26**

(74) Agent: **LECHIEN, Monique**; UCB, S.A., Intellectual Property Department, Allée de la Recherche, 60, B-1070 Bruxelles (BE).

(21) International Application Number: **PCT/EP02/08717**

(22) International Filing Date: **5 August 2002 (05.08.2002)**

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
01119396.8 10 August 2001 (10.08.2001) **EP**

(71) Applicant (*for all designated States except US*): **UCB, S.A.** [BE/BE]; Allée de la Recherche, 60, B-1070 Bruxelles (BE).

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **ATES, Célal** [BE/BE]; Cours d'Orval, 8, B-1348 LOUVAIN-LA-NEUVE (BE). **SURTEES, John** [GB/BE]; Keizerlaan 7, B-3090 JEZUS-EIK (BE). **BURTEAU, Anne-Catherine** [BE/BE]; Rue Béchée 8, B-5031 GRAND-LIEZ (GEMBLoux) (BE). **MARMON, Violeta** [BG/GB]; 26 Kysbie Close, Abingdon-Oxon OX14 1XZ (GB). **CAVOY, Emile** [BE/BE]; Allée de Morfayt, 42, B-6120 HAM-SUR-HEURE (BE).


Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/014080 A2

(54) Title: **OXOPYRROLIDINE COMPOUNDS, PREPARATION OF SAID COMPOUNDS AND THEIR USE IN THE MANUFACTURING OF LLEVETIRACETAM AND ANALOGUES**




(57) Abstract: The present invention relates to an improved process for the preparation of (S)-(-)- α -ethyl-2-oxo-1-pyrrolidine acetamide and analogues thereof. The invention also relates to compounds of the general formula (6) wherein R₁ is methyl or ethyl; and R₂ is C₂#191-C₄#191 alkyl, C₂#191-C₄#191 alkenyl or C₂#191-C₄#191 alkynyl, optionally substituted by one or more halogen, and their preparation processes.

OXOPYRROLIDINE COMPOUNDS, PREPARATION OF SAID COMPOUNDS AND THEIR USE IN THE MANUFACTURING OF LEVETIRACETAM AND ANALOGUES

5 This invention concerns a new and improved process for the preparation of (S)-(-)- α -ethyl-2-oxo-1-pyrrolidine acetamide and analogues thereof, which is referred to under the International Non-proprietary Name of Levetiracetam. Levetiracetam is known as a useful therapeutic agent for the treatment or prevention of epilepsy and other neurological disorders. This invention also discloses novel intermediates and
10 their use in manufacturing processes of Levetiracetam and analogues thereof.

Levetiracetam or (S)-(-)- α -ethyl-2-oxo-1-pyrrolidine acetamide, a laevorotatory compound is disclosed as a protective agent for the treatment and the prevention of hypoxic and ischemic type aggressions of the central nervous system in the European patent No. EP 0 162 036 B and has the following formula.

15




This compound is also effective in the treatment of epilepsy, a therapeutic indication for which it has been demonstrated that its dextrorotatory enantiomer (R)-(-)
20)- α -ethyl-2-oxo-1-pyrrolidine acetamide completely lacks activity (A.J. Gower et al., Eur. J. Pharmacol., 222, 1992, 193-203). A process for the preparation of this dextrorotatory enantiomer has been described in the European patent No. 0165 919.

Manufacturing processes for Levetiracetam have been described in both the European patent No. 0162 036 and in the British patent No. 2 225 322. In the British
25 patent No. 2 225 322 (S)-(-)- α -ethyl-2-oxo-1-pyrrolidine acetamide is prepared by hydrogenolysis of (S)- α -[2-(methylthio)ethyl]-2-oxo-1-pyrrolidineacetamide in the presence of a desulfurizing reagent such as $\text{NaBH}_4/\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$, Raney nickel W-2 or, preferably, Raney nickel T-1. However, this process cannot be conveniently applied on an industrial scale for safety and environmental reasons.

30 Another industrially applicable process was developed and disclosed in a more recent patent application PCT/EP01/01956. The process described in said patent application PCT/EP01/01956 is illustrated in Scheme 1 below. This process is based on the asymmetric hydrogenation of a compound of formula (1), resulting in Levetiracetam (compound of formula (2)). Said patent application also describes the


efficient asymmetric hydrogenation of related compounds of general formula (3), providing the acid and esters of general formula (4).



Scheme 1.

Me represents methyl, and Et represents ethyl.

However, it may be desired to convert the ester (4) directly to Levetiracetam (2) by ammonolysis. A disadvantage of performing said ammonolysis is that racemisation may occur, resulting in the formation of the compound of formula (5) as described in Scheme 2. below.



Scheme 2.


Moreover, the reaction time necessary to obtain a reasonable conversion is generally very long. The reaction time may be decreased by increasing the reaction temperature, but then the extent of racemisation increases to unacceptable levels. No compromise had until now been found between the reaction time, the temperature and extent of racemisation.

It is clear that an industrially viable process without the above-mentioned disadvantage would be extremely desirable.

The process of the present invention largely overcomes the major disadvantages such as the racemisation discussed above and excessive hydrolysis. In addition, the present invention describes novel intermediates and their use in

processes for the preparation of Levetiracetam and analogues thereof. The invention also relates to new processes for preparing said intermediates.

According to a first aspect, the present invention relates to a compound of formula (6):



(6)


wherein R¹ is methyl or ethyl and R² is C₂-C₄ alkyl, C₂-C₄ alkenyl or C₂-C₄ alkynyl, optionally substituted by one or more halogen, preferably F, Cl, Br or I; as well as the stereoisomers and mixtures thereof.

This invention relates to all stereoisomeric forms such as geometrical and optical enantiomeric and diastereoisomeric forms of the compounds of formula (6) and mixtures (including racemates) thereof. The compounds of formula (6) and some of their intermediates have at least one stereogenic center in their structure, being the carbon atom attached to the nitrogen atom of the pyrrolidine heterocycle. This stereogenic center is indicated in formula (6) by an asterisk (*). This stereogenic center may be present in a R or a S configuration, said R and S notation is used in accordance with the rules described in Pure Appl. Chem., 45 (1976) 11-30. The compounds of formula (6) have at least a second stereogenic center in their structure, being the carbon atom of the pyrrolidine cycle to which the R² substituent is attached. This stereogenic center may be in a S or a R configuration. Furthermore certain compounds of formula (6) which contain alkenyl groups may exist as Z or E isomers. In each instance, the invention includes both mixtures and separate individual isomers.

The compound of the formula (6) can be in the form of a solvate, which is included within the scope of the present invention. The solvates are for example hydrates, alcoholates and the like. The compound of the formula (6) can also be in the form of a salt, especially a pharmaceutical acceptable salt, which are also included within the scope of the present invention.

According to a preferred embodiment, the present invention relates to the compound of the general formula (6), wherein the R² substituent is present at position 4 on the ring structure, as given in the following general formula (7) wherein R¹ and R² are as noted above.

4



(7)


According to another preferred embodiment, the present invention relates to the compound of formula (7), wherein the R^2 is a C2-C4 alkyl, C2-C4 alkenyl or C2-C4 alkynyl, optionally substituted by one or more halogen.

5 The term alkyl as used herein includes saturated monovalent hydrocarbon radicals having straight, branched or cyclic moieties or combinations thereof.


The term alkenyl as used herein includes both branched and unbranched unsaturated hydrocarbon radicals having at least one double bond.

10 The term alkynyl as used herein includes both branched and unbranched hydrocarbon radicals having at least one triple bond.


According to a more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is methyl and R^2 is propyl according to the following formula:




15 According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is methyl and R^2 is 2,2-difluorovinyl according to the following formula :




According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is ethyl and R^2 is propyl according to the following formula :




5 According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is ethyl and R^2 is 2,2-difluorovinyl according to the following formula :



10 According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is methyl and R^2 is 2-fluoro-2-methylpropyl according to the following formula :




15 According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is ethyl and R^2 is 2-fluoro-2-methylpropyl according to the following formula :




According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is methyl and R^2 is 2,2-difluoropropyl according to the following formula :

5




According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is ethyl and R^2 is 2,2-difluoropropyl according to the following formula :




10


According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R^1 is methyl and R^2 is cyclopropylmethyl according to the following formula :




According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R¹ is ethyl and R² is cyclopropylmethyl according to the following formula :




According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R¹ is methyl and R² is 2,2,2-trifluoroethyl according to the following formula :



According to yet another more preferred embodiment, the invention relates to the compound of the general formula (7), wherein R¹ is ethyl and R² is 2,2,2-trifluoroethyl according to the following formula :



According to another preferred embodiment, the compound of general formula (6) or (7) is the S isomer as illustrated in the following formula (8) wherein R¹ and R² are as noted above.




In this preferred embodiment the compounds of formula (8) include compounds wherein the second stereogenic center, that is the carbon atom of the pyrrolidine heterocycle to which the R² substituent is attached, is in a S or in a R configuration and their mixtures. Furthermore certain compounds of formula (8) which contain alkenyl groups may exist as Z or E isomers. In each instance, the invention includes both mixtures and separate individual isomers.

The invention also relates to new processes for the manufacture of said compound of the general formula (6) as defined above.


According to a first process, named the "Late Ring-Closure route or LRC route", said compound of general formula (6) of the invention as defined above may be manufactured by a process comprising following steps:

(a) reaction of a compound of formula (9)




with an alcohol of formula R^1OH wherein R^1 is defined as above.

(b) reaction of the corresponding compound of formula (10) thus obtained



with a R^2 -substituted-ethyl-4-bromobutyrate wherein R^2 is defined as above.

(c) cyclisation of the corresponding compound of formula (11) thus obtained



with a catalyst, and


(d) isolation of the resulting compound.

In this process, the compound of formula (9) is an enantiomerically pure or an enantiomerically enriched compound, the chiral centre (either configuration) being denoted by an asterisk (*). By enantiomerically enriched compound is meant a compound containing more than 50 %, preferably more than 55 %, most preferably more than 60 %, of one of the enantiomers. By enantiomerically pure compound is meant a compound containing at least 90 %, preferably at least 95 %, most preferably at least 98 %, of one of the enantiomers.

The first step (step a) of this first process is preferably effectuated in the presence of an alcohol (for instance methanol or ethanol) and thionyl chloride. The second step (step b) is the mono-N-alkylation of the amino-ester of formula (10) with a R^2 -substituted ethyl 4-bromobutyrate (4-EBB) and is preferably effectuated in the presence of an alcohol (for instance methanol, ethanol or isopropanol). The alcohol is preferably isopropanol. The use of isopropanol resulted in a major amount of the monoalkylated ester (11) and a small amount of a dialkylated product which may be separated by column chromatography. Alternatively, the monoalkylated product may be precipitated as its hydrochloride salt by means of gaseous HCl. The hydrochloride of the mono-alkylated product (solid) is next neutralised with aqueous sodium carbonate and extracted with an organic solvent. The second step is preferably performed in the presence of base, most preferably sodium carbonate. The catalyst

used in the third step (step c) in the first process is preferably 2-pyridinol. This reaction is non-racemising and provides enantiomerically enriched or pure (S)-isomers of compounds of formula (8) in the case where the (S) enantiomer of compound (9) is used as starting material.


- 5 According to an alternative process, said compound of general formula (6) of the invention as defined above may be manufactured by a process comprising the step of cyclisation of the compound of formula (11), wherein R^1 and R^2 are as defined above. This process is carried out according to Scheme 4. below :




Scheme 4.

According to a second process, said compounds of formula (6) of the invention as defined above may also be manufactured by a process comprising following steps :

- (a) reaction of an α -ketocarboxylic acid derivative of formula (12)




wherein R^1 is as defined above, with a pyrrolidinone of formula (13)



wherein R^2 is as defined above.

(b) reaction of the corresponding compound of formula (14) thus obtained



(14)


with hydrogen in the presence of an asymmetric hydrogenation catalyst, and

(c) isolation the resulting compound.

This process has as a major advantage that it is much more rapid, simpler, and comprising fewer steps than the first 'LRC' route as discussed above. All details of this process are disclosed in the application PCT/EP01/01956 where it is described for compounds of a more general formula. Said application is hereby further incorporated by reference.


According to a third process, said compounds of the general formula (6) of the invention as defined above may also be manufactured by a process comprising following steps :

(a) reaction of a compound of formula (15)



(15)


wherein R¹ is C₁-C₆ alkyl and X is Cl, Br, I, alkylsulphonate or sulfate; with a pyrrolidone of general formula (13).



(13)

wherein R² is as noted as above;

(b) reaction of the corresponding compound of formula (16) thus obtained






with ethyl-X, wherein X is Cl, Br, I, alkylsulphonate or sulfate and an asymmetric alkylation catalyst or additive;

- 5 (c) optionally, when R¹' is different from R¹, reaction of the compound obtained from step (b) with an alcohol of formula R¹OH, and
 (d) isolating the resulting compound of formula (6).

According to this third process, R¹' is preferably C3-C4 alkyl, especially ter-butyl.

- 10 According to this third process, the asymmetric alkylation catalyst or additive is preferably a chiral amine, most preferably selected from (S)-1-(2-pyrrolidinylmethyl)-pyrrolidine (17), (R)-2-methoxyethoxyethyl-1-phenyl-2-piperidinoethylamine (18) and (S)-1-methyl-2-anilinomethyl pyrrolidine (19).



- 15 Step (b) of this third process is preferably performed in the presence of a base (such as mineral, organic or organometallic bases). The base is preferably butyllithium.

Step (c) of this process is preferably acid or base catalysed.


This process has the advantage that it comprises only few reaction steps.

- 20 Another advantage is that it may be performed using inexpensive and readily available raw materials.


According to a fourth process, the compound of the general formula (6) as defined above may also be prepared by a process comprising following steps :

- (a) reaction of a compound of general formula (20)

13



(20)




(13)

wherein R¹ is as defined above, with a pyrrolidone of general formula (13)

10 wherein R² is defined as above;

(b) separation of the corresponding compound of general formula (21) thus obtained




(21)

wherein R¹ and R² are defined as above;

(c) isolation of the resulting compound of general formula (6).

15 According to this fourth process, the compound of the general formula (6) as defined above may be isolated by industrial chiral chromatographic separation (batch, MCC (Multi Column Chromatography) or SMB (simulated moving bed)) of a compound of general formula (21) according to Scheme 7. below.

14



Scheme 7.

The chromatographic process can be carried out using either the batch or MCC process. Each enantiomer can be separated using a chiral stationary phase to yield enantiomerically pure products.


Commercially available chromatographic columns are for example sold by DAICEL Company or SHISEIDO Company. The preferred DAICEL columns such as the columns sold under the trademark CHIRALPAK AD, CHIRALPAK AS and CHIRALPAK OD were found to be efficient to this end when mobile phases such as mixtures of alkanes with alcohols were used or even a pure alcohol or mixtures of alcohols. The alkane or mixtures of alkanes particularly referred to are: hexane, isohexane or heptane. The alcohol or mixtures of alcohols particularly referred to are: propanol, isopropanol, ethanol or methanol. There is a preference for the use of heptane among the alkanes and there is a preference for the use of ethanol and methanol among the alcohols. There is a preference for the following mixtures: 50% to 95% for the alkane and 50% to 5% for alcohol(s), or 100% of alcohol.

The preferred SHISEIDO columns such as the columns sold under the trademark CERAMOSPHER CHIRAL RU-2 or CERAMOSPHER CHIRAL RU-1 were found to be efficient for the separation when alcohols were used as mobile phase. The alcohols referred to are: propanol, isopropanol, ethanol or methanol. There is a preference for the use of ethanol and methanol among the alcohols.

The extrapolation of small-scale batch separations of this type to an industrial scale proceeds without difficulty in either batch or continuous mode.

According to a second aspect, the present invention also relates to a process for the manufacture of a compound of the general formula (22') wherein $R^{2'}$ is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl or C_2 - C_4 alkynyl, optionally substituted by one or more halogen, said process comprising the ammonolysis of the corresponding compound of formula (6')

15



Scheme 8.

wherein $\text{R}^{1'}$ is $\text{C}_1\text{-C}_6$ alkyl and $\text{R}^{2'}$ is hydrogen, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_2\text{-C}_4$ alkenyl or $\text{C}_2\text{-C}_4$ alkynyl, optionally substituted by one or more halogen, in the presence of water.

Surprisingly, it has been found that performing said ammonolysis in the presence of water greatly overcomes the disadvantages such as racemisation as described in the background art, and encountered when using an organic solvent (e.g. methanol). Other advantage of this invention is minimisation of potential hydrolytic side-reaction.

According to a preferred embodiment, said ammonolysis as described above is performed in a mixture of water and an alcohol. Preferred alcohols are methanol, ethanol, isopropanol and butanol. Most preferably a mixture of water and methanol is used. Using a mixture of water and an alcohol, especially methanol, offers the additional advantage that the level of hydrolysis is even more decreased.

According to a preferred embodiment, said ammonolysis of the invention as described above is performed with NH_3 . Preferably, a 10-95 % (w/w) NH_3 solution in water is used. Most preferably, a 30-80 % (w/w) NH_3 solution in water, especially a 50 % NH_3 solution in water, is used.


According to yet another preferred embodiment, said ammonolysis of the invention as described above is performed at 0 to 40 °C, most preferably at a temperature of 0 to 25°C, especially at a temperature of about 3 to 10 °C.

In the process according to the invention, the molar ratio of NH_3 to the compound of formula (6') is generally at least 1, preferably at least 4, most preferably at least 6. The molar ratio does preferably not exceed 100.

According to a preferred embodiment of the process for the manufacture of the compound of formula (22'), a compound of the general formula (6') is used wherein $\text{R}^{1'}$ is methyl, ethyl or a $\text{C}_3\text{-C}_4$ alkyl. Especially preferred are compounds of general formula (6') wherein $\text{R}^{1'}$ is methyl or ethyl and most preferably wherein $\text{R}^{1'}$ is methyl.


According to another preferred embodiment of the process for the manufacture of the compound of formula (22'), a compound of the general formula (6') is used wherein $\text{R}^{2'}$ is hydrogen.

According to a more preferred embodiment of the process for the manufacture of the compound of formula (22') a compound of the general formula (6') is used wherein R^{1'} is methyl and R^{2'} is hydrogen according to the following formula :



5 The above compound is referred to as PBM (methyl 2-(2-oxo-pyrrolidin-1-yl)butyrate).


According to yet another embodiment of the process for the manufacture of the compound of formula (22'), a compound of the general formula (6') is used wherein R^{1'} is ethyl and R^{2'} is hydrogen according to the following formula :



10


The above compound is referred to as PBE (ethyl 2-(2-oxo-pyrrolidin-1-yl)butyrate).

According to yet another embodiment of the process for the manufacture of the compound of formula (22'), a compound of the general formula (6') is used wherein the R^{2'} substituent is present at position 4 on the ring structure, as given in the following general formula (7') wherein R^{1'} and R^{2'} are as noted above.



20 According to another preferred embodiment of the process according to the invention, the compound of formula (6') is the S isomer as illustrated in the following formula (8') wherein R^{1'} and R^{2'} are as noted above.

17



(8')

The use of an S isomer of formula (8') in the process according to the invention permits to obtain compounds of formula (22') being S isomers. Compounds of formula (6') wherein R^{2'} is different from hydrogen possess a second stereogenic center, being the carbon atom of the pyrrolidine ring to which the R^{2'} substituent is attached. In this case, this stereogenic center may be in an S- or R-form or mixtures of both forms may be used.


According to a more preferred embodiment of the process for the manufacture of the compound of formula (22'), a compound of the general formula (6'), (7') or (8') is used, wherein R^{2'} is selected from the group of hydrogen, propyl, 2,2-difluorvinyl, 2-fluoro-2-methylpropyl, 2,2-difluoropropyl, cyclopropylmethyl and 2,2,2-trifluoroethyl.

The ammonolysis process according to the invention permits high conversion rates. The ammonolysis process according to the invention offers also the advantage that the amount of racemisation and hydrolysis is very low, even negligible. A simple crystallisation of the crude products from this ammonolysis in an organic solvent may give pure compounds, such as pure Levetiracetam.

The compound of formula (6') used as starting material in the process for the manufacture of a compound of formula (22'), can be manufactured by any process suitable therefore.

According to a first variant, the compound of formula (6') is manufactured by a first new process comprising following steps :


(a) reaction of a compound of formula (9)



(9)

with an alcohol of formula R^{1'}OH wherein R^{1'} is defined as above.


(b) reaction of the corresponding compound of formula (10') thus obtained



(10')

with a $R^{2'}$ -substituted-ethyl-4-bromobutyrate wherein $R^{2'}$ is defined as above,

(c) cyclisation of the corresponding compound of formula (11') thus obtained



(11')

in the presence of a catalyst, and

(d) isolation of the resulting compound.


- 15 In this process, the compound of formula (9) is an enantiomerically enriched or an enantiomerically pure compound, the chiral centre (either configuration) being denoted by an asterisk (*).

This first new process as such for the manufacture of a compound of formula (6') is another aspect of the present invention.

- 20 The first step (step a) of this process is preferably performed in the presence of an alcohol (for instance methanol or ethanol) and thionyl chloride. The second step (step b) of this process is the mono-N-alkylation of the amino-ester of formula (10') with a $R^{2'}$ -substituted ethyl 4-bromobutyrate (4-EBB) and is preferably performed in the presence of an alcohol (for instance methanol, ethanol or isopropanol). The alcohol
- 25 is preferably isopropanol. The use of isopropanol presents the further advantage that transesterification did not occur. Moreover, the use of isopropanol resulted in a major amount of the monoalkylated ester (11') and only a small amount of a dialkylated product which may be separated by column chromatography. Alternatively, the monoalkylated product may be precipitated as its hydrochloride salt by means of
- 30 gaseous HCl. The hydrochloride of the mono-alkylated product (solid) is next neutralised with aqueous sodium carbonate and extracted with an organic solvent. The second step is preferably performed in the presence of base, preferably sodium carbonate. The catalyst used in the third step (step c) in the process is preferably 2-pyridinol. This reaction is non-racemising and provides enantiomerically pure (S)-


compounds of formula (8') in the case where the (S) enantiomer of compound (9) is used as starting material.

- According to an alternative process, said compound of general formula (6') of the invention as defined above may be manufactured by a process comprising the step
- 5 of cyclisation of the compound of formula (11'), wherein $R^{1'}$ and $R^{2'}$ are as defined above. This process is carried out according to Scheme 4'. below :




Scheme 4'.

- According to a second variant, the compound of formula (6') is manufactured
- 10 by a second process comprising the following steps :
- (a) reaction of an α -ketocarboxylic acid derivative of formula (12')




wherein $R^{1'}$ is as defined above with a pyrrolidinone of formula (13')



15

wherein $R^{2'}$ is as defined above,

- (b) reaction of the corresponding compound of formula (14') thus obtained



wherein $R^{1'}$ and $R^{2'}$ are defined as above, with hydrogen in the presence of an asymmetric hydrogenation catalyst;


- 5 (c) isolation of the resulting compound.

This second process has as a major advantage that it is much more rapid and simpler, comprising fewer steps than the first 'LRC' route as discussed above. All details of this process are disclosed in the application PCT/EP01/01956 where it is described for compounds of a more general formula. Said application is hereby further


10 incorporated by reference.

According to a third variant, compounds of the general formula (6') as defined above are manufactured by a third new process comprising following steps :

- (a) reaction of a compound of formula (15')




- 15 wherein $R^{1'}$ is as noted above and X is Cl, Br, I, alkylsulphonate or sulfate; with a pyrrolidone of general formula (13')



wherein $R^{2'}$ is as noted as above;

- (b) reaction of the corresponding compound of formula (16') thus obtained




with ethyl-X, wherein X is Cl, Br, I, alkylsulphonate or sulfate in the presence of an asymmetric alkylation catalyst or additive;

- 5 (c) isolation of the resulting compound of formula (6').


According to this third variant, R^{1'} is preferably C₃-C₄ alkyl, especially tert-butyl.

This third new process as such for the manufacture of a compound of formula (6') is another aspect of the present invention.


- 10 According to this third process, the asymmetric alkylation catalyst or additive is preferably a chiral amine, most preferably selected from (S)-1-(2-pyrrolidinylmethyl)-pyrrolidine (17), (R)-2-methoxyethoxyethyl-1-phenyl-2-piperidinoethylamine (18) and (S)-1-methyl-2-anilinomethyl pyrrolidine (19).



(17)



(18)



(19)


15 Step (b) of this process is preferably performed in the presence of a base (such as mineral, organic or organometallic bases). This base is most preferably butyllithium.

- 20 Especially when R^{1'} is not methyl or ethyl, this third process may comprise an additional reaction step wherein the compound obtained from step (b) is reacted with an alcohol of formula R¹OH wherein R¹ is methyl or ethyl, preferably in the presence of an acid, so that a compound of formula (6') is formed wherein R^{1'} is methyl or ethyl.


- 25 This third process has the advantage that it comprises only few reaction steps. Another advantage is that it may be performed using inexpensive and readily available raw materials.

According to a fourth variant, the compound of the general formula (6') as defined above is prepared by a fourth new process comprising following steps :
(a) reaction of a compound of general formula (20')

(a) reaction of a compound of general formula (20')




wherein R¹' is as noted above, with a pyrrolidone of general formula (13')



wherein R²' is defined as above;

(b) separation of the corresponding compound of general formula (21') thus obtained
wherein R¹' and R²' are defined as above; and




15 (c) isolation of the resulting compound of general formula (6').

This fourth new process as such for the manufacture of a compound of formula (6') is another aspect of the present invention.

According to this fourth process, the compound of the general formula (6') as defined above is preferably isolated by industrial chiral chromatographic separation
20 (batch, MCC (Multi Column Chromatography) or SMB (simulated moving bed)) of a compound of general formula (21') according to Scheme 7'. below.

23



Scheme 7'.

According to this fourth process, (S)-PBE and (S)-PBM can be separated using
 5 chiral HPLC by means of commercially available chiral stationary phases. These
 separations can more particularly be performed using chromatographic columns sold
 by DAICEL Company or SHISEIDO Company. The chromatographic process can be
 carried out using either the batch or MCC process. Each enantiomer can be separated
 using a chiral stationary phase to yield enantiomerically pure (S)-PBM and (S)-PBE.

10 The preferred DAICEL columns such as the columns sold under the trademark
 CHIRALPAK AD, CHIRALPAK AS and CHIRALPAK OD were found to be efficient to this
 end when mobile phases such as mixtures of alkanes with alcohols were used or even
 a pure alcohol or mixtures of alcohols. The alkane or mixtures of alkanes particularly
 referred to are: hexane, isohexane or heptane. The alcohol or mixtures of alcohols
 15 particularly referred to are: propanol, isopropanol, ethanol or methanol. There is a
 preference for the use of heptane among the alkanes and there is a preference for the
 use of ethanol and methanol among the alcohols. There is a preference for the
 following mixtures: 50% to 95% for the alkane and 50% to 5% for alcohol(s), or 100%
 of alcohol.

20 The preferred SHISEIDO columns such as the columns sold under the
 trademark CERAMOSPHER CHIRAL RU-2 or CERAMOSPHER CHIRAL RU-1 were
 found to be efficient for the separation when alcohols were used as mobile phase. The
 alcohols referred to are: propanol, isopropanol, ethanol or methanol. There is a
 preference for the use of ethanol and methanol among the alcohols.

25 The extrapolation of small-scale batch separations of this type to an industrial
 scale proceeds without difficulty in either batch or continuous mode.

The optimum conditions as determined by chiral HPLC for the separation of
 both PBE & PBM are presented in Tables I and III below. An estimated productivity for
 PBE and PBM using the MCC process is also given in Tables II and IV.

Table I.

Examples of separation by chiral HPLC : PBM					
Phase provider	Phase	Solvents	k' ₁	Alpha	Resolution
Daicel	Chiralpak® AD	Ethanol 50 % / i-Hexane 50 %	0.499	1.19	1.06
Daicel	Chiralpak® AD	Ethanol 2 % / Methanol 8 % / Hexane 90 %	2.432	1.45	2.1
Daicel	Chiralpak® AD	Acetonitrile 100 %	0.549	1.3	0.79
Daicel	Chiralpak® AD	Ethanol 10 % / Heptane 90 %	3.901	1.24	1.19
Daicel	Chiralpak® AD	Ethanol 5 % / Methanol 5 % Heptane 90 %	3.646	1.41	1.92
Daicel	Chiralpak® AS	i-Propanol 10 % / i-Hexane 90 %	9.408	1.28	2.6
Daicel	Chiralpak® AS	Ethanol 10 % / i-Hexane 90 %	3.035	1.17	1.65
Daicel	Chiralpak® AS	Propanol 10 % / i-Hexane 90 %	2.987	1.14	1.34
Daicel	Chiralpak® OD-H	Ethanol 5 % / i-Hexane 95 %	2.49	1.23	2.97
Daicel	Chiralpak® OD-H	Propanol 5 % / i-Hexane 95 %	1.94	1.22	2.58
Shiseido	Ceramospher Chiral RU-1	Methanol 100 %	4.69	1.28	1.56
Shiseido	Ceramospher Chiral RU-2	Methanol 100 %	3.747	1.29	1.5
Shiseido	Ceramospher Chiral RU-2	Ethanol 100 %	4.853	1.32	1.19

Table II.

Estimated productivity using MCC process : PBM			
Phase provider	Phase	Solvents	Productivity (kg/kg/day)
Daicel	Chiralpak® AD	Ethanol 2 % / Methanol 8 % / i-Hexane 90 %	0.17

5 Productivity as presented in the above table is expressed as Kg of racemic PBM engaged per Kg of chiral stationary phase per day.

Table III.

Examples of separation by chiral HPLC : PBE					
Phase provider	Phase	Solvents	k' ₁	Alpha	Resolution
Daicel	Chiralpak® AD	Ethanol 50 % / i-Hexane 50 %	0.449	1.3	1.15
Daicel	Chiralpak® AD	Ethanol 2 % / Methanol 8 % / Hexane 90 %	1.955	1.9	3.32
Daicel	Chiralpak® AD	Acetonitrile 100 %	0.554	1.8	2.05
Daicel	Chiralpak® AD	Ethanol 10 % / Heptane 90 %	3.076	1.5	4.4
Daicel	Chiralpak® AD	Ethanol 5 % / Methanol 5 % Heptane 90 %	2.971	1.7	2.93
Daicel	Chiralpak® AD	Methanol 5 % / Benzine 95 %	3.227	1.7	2.99
Daicel	Chiralpak® AD	i-Propanol 10 % / i-Hexane 90 %	5.029	2.16	7.39
Daicel	Chiralpak® AD	Ethanol 10 % / i-Hexane 90 %	1.764	1.9	5.97
Daicel	Chiralpak® AD	Propanol 10 % / i-Hexane 90 %	1.733	1.86	5.46
Daicel	Chiralpak® AD	Ethanol 5 % / i-Hexane 95 %	1.878	1.13	1.66
Daicel	Chiralpak® AD	Propanol 5 % / i-Hexane 95 %	1.44	1.14	1.56
Shiseido	Ceramospher Chiral Ru-1	Methanol 100 %	5.047	1.89	3.57
Shiseido	Ceramospher Chiral Ru-2	Methanol 100 %	3.869	1.84	3.21
Shiseido	Ceramospher Chiral Ru-2	Ethanol 100 %	3.97	2.01	1.94

10 Table IV.

Estimated productivity using MCC process : PBE			
Phase provider	Phase	Solvents	Productivity (kg/kg/day)
Daicel	Chiralpak® AD	Ethanol 10 % / Heptane 90 %	0.84

Productivity as presented in the above table is expressed as Kg of racemic PBE engaged per Kg of chiral stationary phase per day.

In the implementation of the processes according to the invention, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example extraction,
5 crystallisation, distillation and chromatography, or any combination of the same.

Stereoisomerically pure forms of said compounds of the invention (and said intermediates) can be obtained by the application of procedures known to a chemist skilled in the art. For example, diastereoisomers can be separated by physical
10 methods such as selective crystallisation or chromatographic techniques, e.g. counter current distribution, liquid chromatography and related methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereoisomeric salts or compounds; then physically separating said mixtures of
15 diastereoisomeric salts or compounds by, for example, selective crystallisation or chromatographic techniques, e.g. liquid chromatography and related methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers.

Alternatively, pure stereochemically isomeric forms may be obtained by using
20 enantioselective reactions according to procedures known by the person skilled in the art.


Another alternative manner of separating the enantiomeric forms of the compounds of formula (6) or (6') and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

According to another aspect, the present invention also relates to any
25 compounds obtained by a process of the invention as defined above. In particular, the invention comprises Levetiracetam obtained by said processes. More particularly, the present invention also relates to new compounds obtainable by the processes according to the invention such as compounds of formula (22') wherein R^{2'} is 2-fluoro-
30 2-methylpropyl or cyclopropylmethyl. More specifically the present invention also relates to the (4S) and (4R) diastereoisomers of (2S)-2-[4-(2-fluoro-2-methylpropyl)-2-oxo-1-pyrrolidinyl]butanamide and of (2S)-2-[4-cyclopropylmethyl)-2-oxo-1-pyrrolidinyl]butanamide, and pharmaceutical compositions containing such compounds and their use as pharmaceuticals.

35 The following examples serve to illustrate the invention and therefore should not be taken to limit the scope thereof.

EXAMPLESExample 1

Step 1 - Synthesis of methyl (S)-aminobutyrate hydrochloride



5.0g of (S)-amino butyric acid (23) was suspended in 50 ml of methanol and stirred at 0-5°C. 6.35g of thionyl chloride was added dropwise over 45 min to form a clear solution. After stirring for 20 hours at room temperature, the reaction was concentrated under reduced pressure to dryness and the almost colourless residue solidified to give the required product which was dried in an oven at 50°C under vacuum (7.6g; 102% crude yield). The same reaction was scaled-up from 200g of the amino acid and provided 296g (99.5% yield) of product (24).

Analysis gave the following results:

¹H NMR (DMSO-d₆) : δ 0.94 (3H, t) 1.88 (2H, q) 3.75 (3H, s) 3.9 (1H, m) 8.8 (3H, m).

m.p. : 107°C-110°C


IR : 2876 cm⁻¹, 1742 cm⁻¹.

TLC : SiO₂, 20%MeOH/80%EtOAc/1%NH₄OH, UV & IR.

(TLC is an abbreviation for thin layer chromatography).

Step 2 - Synthesis of methyl (S)-aminobutyrate-N(4-ethylbutyrate)

27



Scheme 10.

2.0 g of (S)-aminobutyrate hydrochloride salt (24) was dissolved and stirred at room temperature in 20 ml of 2-propanol, followed by addition of 2.8g of sodium carbonate and the reaction was then heated to reflux. When reflux temperature was reached, 2.8g of 4-BBE (ethyl-4-bromobutyrate) was added dropwise over a period of 10 min, with reflux and stirring being maintained for 24 hrs. The reaction medium was allowed to cool to room temperature, the salts were filtered and rinsed with 50 ml of 2-propanol. Following this alkylation the desired product (25) may be isolated and purified either by chromatography or via the hydrochloride salt (25') as depicted in Scheme 10. above and as described in Methods A and B below.

(Method A) : The filtrate was concentrated under reduced pressure to give 3.0g of a pale yellow liquid. This liquid was purified by chromatography through 125g of silica and eluted with a 50/50 mixture of hexane/ethyl acetate to provide the required 2.45g (81% yield) mono alkylated ester (25) (Method B): Chromatography can be avoided if the corresponding hydrochloride salt is generated, precipitated and filtered from a mixture of isopropanol and DIPE (di-isopropylether). Treatment of this salt (25') with sodium carbonate in water and extraction with ethyl acetate and concentration provides the pure free base (25) (the required mono alkylated ester) as a liquid.

Analysis gave the following results:


¹H NMR (CDCl₃) : δ 0.9 (3H, t) 1.2 (2H, t) 1.4 (1H, s) 1.5-1.7 (4H, m) 2.3-2.7 (4H, m) 3.15 (1H, t) 3.7 (3H, s) 4.1 (2H, q).

The identity of the product is confirmed by GC-MS, TLC.

IR : 2938 cm^{-1} , 1730 cm^{-1} .

TLC : SiO_2 , 50%Hexane/50%EtOAc, UV & IR.


Step 3 - Synthesis of methyl (S)-pyrrolidino-butyrates (26) [(S)-PBM]



1.0g of compound (25) and 2-pyridinol (0.02g; 5 mol%) were magnetically stirred in 5 ml of toluene at reflux for 24 hrs. The reaction mixture was allowed to cool to room temperature and TLC analysis showed almost complete conversion. The reaction mixture was then evaporated under reduced pressure to leave crude (S)-PBM (26) as a pale brown liquid (1.0g).


The identity of the product was confirmed by GC-MS, TLC, HPLC (Chiral and Achiral) using external references.

15 Step 4 - Ammonolysis of (S)-PBM to give Levetiracetam.



11.3g of ammonia gas was condensed in 13.2 ml of water at approximately 0°C and the temperature was maintained at 0-5°C. Then 20g of (S)-PBM (26) was added dropwise over a period of 10 min and reaction mixture was maintained at 5°C and stirred for minimum 8 hrs (reaction was complete as indicated by TLC). The reaction mixture was then evaporated to dryness under vacuum and dried by means of toluene (2x50 ml) to give minimum 17g (92%) of crude (S)-pyrrolidinobutyramide (crude Levetiracetam) as an off-white to beige solid.

25 Analysis gave the following results (chiral and achiral HPLC): The extent of racemisation was 0.0%. The extent of hydrolysis was measured to 2.5%.


Example 2

5 17.3g of ammonia gas were condensed in 22 ml of water at 0°C and temperature maintained at 0-5°C. Then 20g of (S)-PBE obtained via SMB separation of the corresponding racemic mixture were added dropwise over a period of 2 min and the reaction mixture was maintained at 5°C and stirred for 96 hrs (reaction was complete as judged by TLC). The reaction mixture was then evaporated to dryness
10 under vacuum and dried by means of toluene (2x100 ml) to give minimum 14.8g (87%) of crude (S)-pyrrolidinobutyramide as a brown orange solid. Analysis gave the following results (chiral and achiral HPLC): The extent of racemisation was 1.6% with 6.6% hydrolysis.

15 Example 3

 10.3g of ammonia gas were condensed in 13.2 ml water at 0°C and the temperature of the system was maintained at 0-5°C. 20g of (S)-PBE obtained via asymmetric hydrogenation was then added dropwise over a period of 10 min, maintaining the reaction mixture at 5°C. The system was then stirred for 96 hrs, with
20 TLC indicating completion of reaction. The reaction mixture was then evaporated to dryness under vacuum and dried by means of toluene (2x50 ml) to give minimum 15.7g (92%) of crude (S)-pyrrolidinobutyramide as a brown orange solid. Analyses gave the following results (chiral and achiral HPLC): The extent of racemisation was 0.2% with 3.4% hydrolysis.

Example 4.



A reaction flask was charged with the chiral amine (34) (1.07 equivalent (eq.);
 5 and anhydrous toluene (15vol) with stirring under an inert atmosphere. The solution
 was cooled below -70°C and BuLi (2.5M in hexane, 1.04 eq.) was added dropwise. The
 reaction mixture was stirred for 30 min at this temperature, then at -10° to 0°C for 10
 min. A solution of t-butyl 2-(2-oxopyrrolidin-1-yl)-acetate (32) (600mg, 1 eq., 1wt) in
 toluene (5 vol) was added slowly, keeping the reaction temperature below -70°C . The
 10 reaction mixture was stirred at -40 to -50°C for 30 min. Ethyl iodide 2.5eq., 1vol) was
 then added and the reaction mixture was stirred at -50 to -40°C for 3 hrs. After being
 kept in the freezer at approximately -40°C overnight, the reaction mixture was diluted
 with pH 7 buffer ($\text{KH}_2\text{PO}_4/\text{KOH}$, 1M, 33vol) and dichloromethane (33vol). The aqueous
 phase was extracted with dichloromethane (3 x 16vol) and the combined organic
 15 extracts were then dried over MgSO_4 and concentrated *in vacuo* to give crude material.
 Purification of this crude product using flash chromatography (SiO_2 , 40wt) with
 hexane/EtOAc eluent gave the desired alkylated product (33) in 78% yield.

$^1\text{H-NMR}$ in CDCl_3 : δ 0.85t(3H), 1.4s(9H), 1.5-1.7m(1H), 1.9-2.0m(3H), 2.45m(2H),
 20 3.25m (1H), 3.5m(1H), 4.5dd(1H)

HPLC analysis : t-Butyl 2-(2-oxopyrrolidin-1-yl)-butanoate (25mg) was accurately
 weighed into a 25ml volumetric flask. Mobile phase (99:1 hexane/isopropanol, 20ml,) was added and the sample was dissolved using ultrasonication. After cooling to
 ambient temperature the concentration was adjusted with mobile phase to give a
 25 working concentration of 1mg/ml. The analysis was conducted using a column sold
 under the trademark CHIRACEL OD (4.6 x 250mm, DAICEL), flow rate of 1ml/min,
 UV detection at 250nm and injection volume of 20 μl at ambient temperature. The
 relative retention times of the two enantiomers was 17.9 and 22.3 minutes

TLC conditions: SiO₂ in EtOAc; visualisation with KMnO₄.

Example 5:

1. Evaluation of type of solvent most suitable for ammonolysis of (S)-PBE.

5 The ammonolysis of (S)-PBE was investigated in the presence of water, toluene, methanol and ethyl acetate. It was shown that the ammonolysis of (S)-PBE can only be successfully realized in the presence of water. When using methanol, the reaction is very slow and when using the other solvents mentioned above the extent of reaction is minimal.

10 2. Evaluation of optimum reaction temperature for the ammonolysis of (S)-PBE to form Levetiracetam.

15 The ammonolysis of (S)-PBE was carried out either at room temperature or at 40°C using (S)-PBE (1 equivalent) in the presence of water (6,5 volume) and various concentrations of NH₃ (15, 10, 7, 5, and 2 equivalents). The reactions were carried out at room temperature and 40°C, being followed by TLC for at least 24 hours. At the end of the reaction the extent of racemisation and hydrolysis was determined by HPLC.

It was shown that :

- 20 - good conversion was obtained, especially when at least 4 equivalents of NH₃ (per eq. of (S)-PBE) were used;
- the extent of racemisation did not exceed 8% at 40 °C and decreased with reaction temperature. At temperatures between 0 and 25 °C, the extent of racemisation was less than 3%;
- 25 - the amount of hydrolysis was low, especially at higher molar ratios of NH₃ to (S)-PBE.

3. Evaluation of different concentrations of NH₃ for ammonolysis of (S)-PBE.

30 Six experiments were performed in a 100ml reactor while varying the concentration of NH₃ and reaction temperature. (S)-PBE (1 equivalent) was mixed with 10 equivalents of NH₃ from either a commercial solution of NH₃ (28% w/w) or a more concentrated solution (± 50 % w/w). The temperatures used were either 5, 10 or 20°C. The reaction was followed by TLC until no (S)-PBE remained and the extent of hydrolysis and racemisation was determined by HPLC.

It was shown that :

- 35 - a more concentrated solution of NH₃ did not substantially influence the extent of racemisation,
- the extent of racemisation was always less than 3% at all reaction temperatures which were tested,
- the extent of racemisation increases only very moderately between 5 and 20 °C.

- the extent of hydrolysis was low, especially when using concentrated NH_3 solution ($\pm 50\%$ w/w),
- the extent of racemisation is always lower at lower reaction temperature.

In summary, the following conclusions can be made:

- 5 - the ammonolysis can easily be performed in the presence of water (containing preferably at least 4 equivalents of NH_3), this reaction does not require any catalyst and may be performed in less than 24 hours,
- the extent of racemisation is low (less than 3% when reaction temperature is less than 20°C), and concentration of NH_3 was found to have only a minor influence on
- 10 the racemisation,
- the extent of hydrolysis can be reduced in an even more substantial way when using a more concentrated solution of NH_3 ($\pm 50\%$ w/w) at low reaction temperature (reaction takes less than 48 hours).

15 Example 6 :

(S)-PBE was reacted under the conditions specified in Table VI. The results are summarised in Table VI. below.




Table VI.

N° Exp.	Reaction conditions					HPLC Analysis area %		
	(S)- PBE (g.)	NH_3 (eq.)	H_2O (Vol.)	Time (hrs)	T° (°C)	acid (% area)	Levetiracetam or (S)-Amide (% area)	(R)- Amide (% area)
6	20	6.2	0.66	96h00	5	3.44	97.85	2.00

The starting material contained 1.6 % of the (R)-enantiomer and 98.4 % of the (S)-enantiomer. The difference in enantiomeric purity between the starting material and the final amides obtained was 0.4 %. This result corresponds to the degree of racemisation accompanied by said ammonolysis.

The product obtained from the experiment described above was recrystallised in eight volumes of acetone and filtered at 2°C . to give the final product, (S)-(-)- α -ethyl-

2-oxo-1-pyrrolidine acetamide or Levetiracetam in 69.1% yield. The recrystallised product contained 0.11 % of the (R)-amide product and 0.08 % of hydrolysed product.

Example 7 :

(S)-PBM was reacted under the conditions specified in Table VIII. The results are summarised in Table VIII. below.




Table VIII.


N° Exp.	Reaction conditions					HPLC Analysis area %		
	(S)- PBM (g)	NH ₃ (eq.)	H ₂ O (Vol.)	Time (hrs)	T° (°C)	acid (% area)	Levetiracetam (S)-amide (% area)	Opposite (R)-amide (% area)
22	22	6.0	0.66	16h40	5	3.68	96.31	2.53

The starting material contained 96.3 % of the (S)-enantiomer and 3.5 % of the (R)-enantiomer. The difference in enantiomeric purity between the final product Levetiracetam and the starting material (S)-PBM was approximately 0.2 %, indicating indeed that the ammonolysis is accompanied by a negligible racemisation in this case.

The final product obtained from the experiment above was recrystallized from eight volumes of acetone and filtered at 4°C. Levetiracetam is obtained in 73.3 % yield. The recrystallized product contained 1.64% of the opposite enantiomeric amide and 0.03 % of the hydrolysed product. Recrystallisation in the presence of acetone as described allows production of Levetiracetam of a sufficient quality for commercial purposes.

The same reaction was finally performed on an increased scale according to Scheme 18. below. Racemisation was as previously observed negligible (0.2 %) .

34



Scheme 18.

5 In summary, it has been shown that Levetiracetam may be obtained via ammonolysis of (S)-PBE in concentrated NH_3 (50% in water) and at 5°C. Scaling-up of this reaction has been successfully demonstrated in 0.6 volumes of water in the presence of 6 equivalents of NH_3 . The extent of racemisation varies between 0.4 and 2.0%, that of hydrolysis between 3.5 and 6.6 %, with a reaction time of approximately 10 96 hours.

Alternatively, Levetiracetam may equally be obtained via ammonolysis of (S)-PBM in 0.6 volumes of water containing 6 equivalents of NH_3 and at 5°C. The reaction time is much shorter and can be realised in 8 to 10 hours. The extent of racemisation varies between 0.0 and 0.2 % and that of hydrolysis ranges from 1.8 to 3.6 %.

15 Example 8

8.1 Preparation of methyl (2S)-2-[2-oxo-(4S)-4-propyl-1-pyrrolidinyl]butanoate

A reaction flask was charged with 2 g of methyl (Z)-2-[2-oxo-(4S)-4-propyl-1-pyrrolidinyl]-2-butenate, 20 ml of anhydrous and degassed methanol and 27 mg of (S,S)-Me-DUPHOS/ $\text{Rh}(\text{BF}_4)$. The reaction flask was purged with hydrogen and the 20 hydrogen pressure was adjusted to 10 atm. This reaction mixture was stirred during about 20 hours at room temperature and then concentrated. 1.96 g of methyl (2S)-2-[2-oxo-(4S)-4-propyl-1-pyrrolidinyl]butanoate was obtained.


8.2 Ammonolysis

Ammonia gas was condensed in 2 ml water at 0-5°C and the temperature of 25 the system was maintained at 0-5°C. 0.68 g of methyl (2S)-2-[2-oxo-(4S)-4-propyl-1-

- pyrrolidinyl]butanoate obtained such as described above was then added dropwise, maintaining the reaction mixture at 0-5°C. The system was then stirred for 6 hrs. with TLC indicating completion of reaction. After standing overnight at ambient temperature the reaction mixture was concentrated at 40°C under vacuum and
- 5 further dried by azeotropic distillation with toluene to give 150 mg of crude (2S)-2-[2-oxo-(4S)-4-propyl-1-pyrrolidinyl]butanamide.

CLAIMS

1. A compound of formula (6)



(6)


wherein

R¹ is methyl or ethyl ; and

R² is C₂-C₄ alkyl, C₂-C₄ alkenyl or C₂-C₄ alkynyl, optionally substituted by one or more halogen;

as well as the stereoisomers and mixtures thereof.

2. The compound according to claim 1, wherein the R² substituent is present at position 4 on the ring structure, according to the following general formula (7).




(7)

3. The compound according to claim 2, wherein R¹ is methyl and R² is propyl or 2,2-difluorovinyl.
4. The compound according to claim 2, wherein R¹ is ethyl and R² is propyl or 2,2-difluorovinyl.
5. The compound according to claim 2, wherein R¹ is methyl and R² is a substituent selected from 2-fluoro-2-methylpropyl, 2,2-difluoropropyl, cyclopropylmethyl and 2,2,2-trifluoroethyl.

6. The compound according to claim 2, wherein R^1 is ethyl and R^2 is a substituent selected from 2-fluoro-2-methylpropyl, 2,2-difluoropropyl, cyclopropylmethyl and 2,2,2-trifluoroethyl.

5


7. The compound according to any of claims 1 to 6, which is an S isomer, according to the following formula (8)



(8)

8. A process for the manufacture of a compound according to any of claims 1 to 7, said process comprising following steps :
(a) reaction of a compound of formula (9)

15




(9)

with an alcohol of formula R^1OH wherein R^1 is as noted in claim 1,


20

- (b) reaction of the corresponding compound of formula (10) thus obtained



(10)

- with a R^2 -substituted-ethyl-4-bromobutyrate wherein R^2 is as noted in claim 1,
(c) cyclisation of the corresponding compound of formula (11) thus obtained




(11)

with a catalyst,

(d) isolation of the resulting compound.


- 5 9. The process according to claim 8, wherein step (a) is performed in the presence of thionyl chloride and an alcohol.
- 10 10. The process according to claims 8 or 9, wherein step (b) is performed in the presence of a base and an alcohol.
- 10 11. The process according to any of claims 8 to 10, wherein the catalyst used in step (c) is pyridinol.
12. The process for the manufacture of a compound according to any of claims 1 to 7,
 - 15 said process comprising a step of cyclisation of the compound of formula (11)



(11)


wherein R¹ and R² are as in claim 1:

13. A process for the manufacture of a compound according to any of claims 1 to 7,
 - 20 said process comprising following steps :
 - (a) reaction of an α -ketocarboxylic acid derivative of formula (12)




(12)

wherein R^1 is as noted in claim 1, with a pyrrolidinone of formula (13)



(13)

- 5 wherein R^2 is as noted in claim 1,
 (b) reaction of the corresponding compound of formula (14) thus obtained




(14)

- with hydrogen in the presence of an asymmetric hydrogenation catalyst, and
 10 (c) isolation of the resulting compound.

14. A process for the manufacture of a compound according to any of claims 1 to 7,
 said process comprising following steps :
 (a) reaction of a compound of formula (15)


15

40



(15)

wherein R¹ is C₁-C₆ alkyl and X is Cl, Br, I, alkylsulphonate or sulfate; with a pyrrolidone of formula (13)




(13)

5

wherein R² is as in claim 1;

(b) reaction of the corresponding compound of formula (16) thus obtained



(16)

with ethyl-X, wherein X is Cl, Br, I, alkylsulphonate or sulfate in the presence of an asymmetric alkylation catalyst or additive;


10

(c) optionally, when R¹ is different from R¹, reaction of the compound obtained in step (b) with an alcohol of formula R¹OH, and
(d) isolation of the resulting compound of formula (6).

15 15. A process for the manufacture of a compound according to any of claims 1 to 7, said process comprising following steps :


(a) reaction of a compound of general formula (20)

41



(20)


wherein R^1 is as defined in claim 1, with a pyrrolidone of general formula (13)



(13)

5 wherein R^2 is defined as in claim 1;

(b) separation of the corresponding compound of formula (21) thus obtained




(21)

wherein R^1 and R^2 are defined as in claim 1;


(c) isolation of the resulting compound of formula (6).

10 16. A process for the manufacture of a compound of formula (22')



(22')

wherein $R^{2'}$ is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl or C_2 - C_4 alkynyl, optionally substituted by one or more halogen, said process comprising the ammonolysis of the corresponding compound of formula (6')




(6')

wherein $R^{1'}$ is C_1 - C_6 alkyl and $R^{2'}$ is hydrogen, C_1 - C_4 alkyl, C_2 - C_4 alkenyl or C_2 - C_4 alkynyl, optionally substituted by one or more halogen, in the presence of water.

- 10 17. The process according to claim 16, wherein said ammonolysis is performed in a mixture of water and an alcohol.
18. The process according to claim 16 or 17, wherein said ammonolysis is performed in a 30-80% (w/w) NH_3 solution in water.
- 15 19. The process according to any of claims 16 to 18, wherein said ammonolysis is performed at 0 to 25°C.
20. The process according to any of claims 16 to 19, wherein the molar ratio of NH_3 to the compound of formula (6') is at least 4.
- 20 21. The process according to any of claims 16 to 20, wherein a compound of formula (6') is used wherein $R^{1'}$ is methyl and $R^{2'}$ is hydrogen.
- 25 22. The process according to any of claims 16 to 20, wherein a compound of formula (6') is used wherein $R^{1'}$ is ethyl and $R^{2'}$ is hydrogen.
23. The process according to any of claims 16 to 20, wherein a compound of formula (6') is used wherein the $R^{2'}$ substituent is present at position 4 on the ring structure, according to the following general formula (7')
- 30


43



(7')


24. The process according to any of claims 16 to 20 or 23, wherein a compound of formula (6') or (7') is used wherein $R^{2'}$ is selected from the group of propyl, 2,2-difluorovinyl, 2-fluoro-2-methylpropyl, 2,2-difluoropropyl, cyclopropylmethyl and 2,2,2-trifluoroethyl.

25. The process according to any of claims 16 to 24, wherein compound (6') is an S isomer according to the following formula (8')



(8')


26. The process according to any of claims 16 to 25, wherein compound (6') is obtained by a process comprising following steps :
(a) reaction of a compound of formula (9)



(9)

with an alcohol of formula $R^{1'}OH$ wherein $R^{1'}$ is as noted in claim 16,

(b) reaction of the corresponding compound of formula (10') thus obtained




(10')

with a R^{2'}-substituted-ethyl-4-bromobutyrate wherein R^{2'} is as noted in claim 16,

(c) cyclisation of the corresponding compound of formula (11') thus obtained

5




(11')

in the presence of a catalyst, and

(d) isolation of the resulting compound.

27. The process according to any of claims 16 to 25, wherein compound (6') is

10 obtained by a process comprising a step of cyclisation of a compound of formula (11')




(11')

wherein R^{1'} and R^{2'} are as noted in claim 16.


28. The process according to any of claims 16 to 25, wherein compound (6') is obtained by a process comprising following steps :

(a) reaction of an α -ketocarboxylic acid derivative of formula (12')



(12')

5 wherein R¹' is as noted in claim 16, with a pyrrolidinone of formula (13')




(13')

wherein R²' is as noted in claim 16.

(b) reaction of the corresponding compound of formula (14') thus obtained

10




(14')

with hydrogen in the presence of an asymmetric hydrogenation catalyst, and
(c) isolation of the resulting compound.

15 29. The process according to any of claims 16 to 25, wherein compound (6') is obtained by a process comprising following steps :


(a) reaction of a compound of formula (15')

46



(15')

wherein R^{1'} is as noted in claim 16 and X is Cl, Br, I, alkylsulphonate or sulfate;
with a pyrrolidone of formula (13')




(13')

5

wherein R^{2'} is as in claim 16,

(b) reaction of the corresponding compound of formula (16') thus obtained



(16')


with ethyl-X, wherein X is Cl, Br, I, alkylsulphonate or sulfate in the presence of
an asymmetric alkylation catalyst or additive,

10

(c) isolation of the resulting compound.


30. The process according to any of claims 16 to 25, wherein compound (6') is
obtained by a process comprising following steps :

(a) reaction of a compound of general formula (20')




(20')

wherein $R^{1'}$ is as noted in claim 16, with a pyrrolidone of general formula (13')



wherein $R^{2'}$ is defined as in claim 16.

5 (b) separation of the corresponding compound of formula (21') thus obtained, and



(c) isolation of the resulting compound of formula (6').

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 February 2003 (20.02.2003)

PCT

(10) International Publication Number
WO 03/014080 A3

- (51) International Patent Classification⁷: C07D 207/26
- (74) Agent: LECHIEN, Monique; UCB, S.A., Intellectual Property Department, Allée de la Recherche, 60, B-1070 Bruxelles (BE).
- (21) International Application Number: PCT/EP02/08717
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 5 August 2002 (05.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
01119396.8 10 August 2001 (10.08.2001) EP
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): UCB, S.A. [BE/BE]; Allée de la Recherche, 60, B-1070 Bruxelles (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): ATEs, Célal [BE/BE]; Cours d'Orval, 8, B-1348 LOUVAIN-LA-NEUVE (BE). SURTEES, John [GB/BE]; Keizerlaan 7, B-3090 JEZUS-EIK (BE). BURTEAU, Anne-Catherine [BE/BE]; Rue Béchée 8, B-5031 GRAND-LEEZ (GEMBLOUX) (BE). MARMON, Violeta [BG/GB]; 26 Kysbie Close, Abingdon-Oxon OX14 1XZ (GB). CAVOY, Emile [BE/BE]; Allée de Morfayt, 42, B-6120 HAM-SUR-HEURE (BE).
- Published:
— with international search report
- (88) Date of publication of the international search report:
6 November 2003
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: OXOPYRROLIDINE COMPOUNDS, PREPARATION OF SAID COMPOUNDS AND THEIR USE IN THE MANUFACTURING OF LEVETIRACETAM AND ANALOGUES

(57) Abstract: The present invention relates to an improved process for the preparation of (S)-(-)- α -ethyl-2-oxo-1-pyrrolidine acetamide and analogues thereof. The invention also relates to compounds of the general formula (6) wherein R₁ is methyl or ethyl; and R₂ is C²-C⁴191 alkyl, C²191-C⁴191 alkenyl or C²191-C⁴191 alkynyl, optionally substituted by one or more halogen, and their preparation processes.

WO 03/014080 A3

INTERNATIONAL SEARCH REPORT

Inter national Application No
PCT/EP 02/08717A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D207/26

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)
EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 1 309 692 A (UCB SA) 14 March 1973 (1973-03-14) cited in the application page 3	1,2,16
A	EP 0 165 919 A (UCB SA) 27 December 1985 (1985-12-27) cited in the application the whole document	1-30
A	EP 0 162 036 A (UCB SA) 21 November 1985 (1985-11-21) cited in the application the whole document	1-30
-/--		

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

12 August 2003

Date of mailing of the international search report

21/08/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Timmermans, M

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 02/08717

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 64637 A (DIFFERDING EDMOND ; SURTEES JOHN (BE); UCB FARCHIM S A AG LTD (CH);) 7 September 2001 (2001-09-07) cited in the application page 6-8 -----	1,2,4,13

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08717

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 1309692	A	14-03-1973	BE 762728 A4	10-08-1971
			CH 530395 A	15-11-1972
			CS 167911 B2	28-05-1976
			CS 167912 B2	28-05-1976
			CY 727 A	01-03-1974
			DE 2106418 A1	19-08-1971
			DK 137451 B	06-03-1978
			ES 388135 A2	01-05-1973
			FI 54916 B	29-12-1978
			FR 2081508 A6	03-12-1971
			JP 53005301 B	25-02-1978
			JP 986288 C	07-02-1980
			JP 53130656 A	14-11-1978
			JP 54020491 B	23-07-1979
			MY 7274 A	31-12-1974
			NL 7101697 A ,B,	17-08-1971
			SU 508185 A3	25-03-1976
			SU 508186 A3	25-03-1976
			YU 30671 A ,B	31-10-1979
			YU 144578 A1	31-10-1982
			ZA 7100877 A	27-10-1971
EP 0165919	A	27-12-1985	AT 45348 T	15-08-1989
			AU 574175 B2	30-06-1988
			AU 4252985 A	21-11-1985
			BG 50272 A3	15-06-1992
			BG 60030 A3	15-07-1993
			CA 1237138 A1	24-05-1988
			CN 85105307 A ,B	14-01-1987
			CY 1568 A	20-12-1991
			DE 3572168 D1	14-09-1989
			DK 212885 A ,B,	16-11-1985
			EP 0165919 A1	27-12-1985
			ES 8703419 A1	01-05-1987
			ES 8706114 A1	16-08-1987
			FI 851876 A ,B,	16-11-1985
			GR 851156 A1	25-11-1986
			HK 62692 A	28-08-1992
			IE 58393 B1	08-09-1993
			IL 75180 A	31-05-1988
			JP 60255764 A	17-12-1985
			KR 9203818 B1	15-05-1992
			NO 851934 A ,B,	18-11-1985
			PL 253373 A1	06-05-1986
			PL 257386 A1	07-10-1986
			PT 80461 A ,B	01-06-1985
			SG 80190 G	23-11-1990
			SU 1417799 A3	15-08-1988
			SU 1320207 A1	30-06-1987
			SU 1428194 A3	30-09-1988
			US 4837224 A	06-06-1989
			US 4696942 A	29-09-1987
			ZA 8503636 A	24-12-1985
EP 0162036	A	21-11-1985	AT 45567 T	15-09-1989
			AU 574465 B2	07-07-1988
			AU 4253085 A	20-11-1986
			BG 47497 A3	16-07-1990

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 02/08717

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0162036	A		BG 50156 A3	15-05-1992
		CA 1235129 A1	12-04-1988	
		CN 85105301 A ,B	14-01-1987	
		CY 1567 A	20-12-1991	
		DE 3572348 D1	21-09-1989	
		DK 212985 A ,B,	16-11-1985	
		EP 0162036 A1	21-11-1985	
		ES 8608485 A1	01-12-1986	
		ES 8704893 A1	01-07-1987	
		FI 851875 A ,B,	16-11-1985	
		GR 851155 A1	25-11-1985	
		HK 52391 A	19-07-1991	
		IE 59950 B1	04-05-1994	
		IL 75179 A	31-05-1988	
		JP 1901367 C	27-01-1995	
		JP 6029186 B	20-04-1994	
		JP 60252461 A	13-12-1985	
		KR 9203818 B1	15-05-1992	
		LT 2584 R3	25-03-1994	
		LU 90615 A9	02-10-2000	
		LU 90682 A9	30-01-2001	
		LV 5233 A3	10-10-1993	
		NO 851933 A ,B,	18-11-1985	
		PL 253374 A1	06-05-1986	
		PL 257385 A1	07-10-1986	
		PT 80460 A ,B	01-06-1985	
		SG 80090 G	23-11-1990	
		SU 1402260 A3	07-06-1988	
		SU 1430392 A1	15-10-1988	
		SU 1428195 A3	30-09-1988	
		US 4837223 A	06-06-1989	
		US 4943639 A	24-07-1990	
		US 4696943 A	29-09-1987	
		ZA 8503635 A	24-12-1985	
WO 0164637	A	07-09-2001	AU 5214401 A	03-09-2001
			AU 7389601 A	12-09-2001
			BG 107004 A	30-04-2003
			BG 107016 A	30-04-2003
			BR 0108657 A	29-04-2003
			BR 0108664 A	29-04-2003
			CA 2401033 A1	30-08-2001
			CA 2401048 A1	07-09-2001
			CN 1404469 T	19-03-2003
			CN 1404470 T	19-03-2003
			CZ 20022849 A3	12-02-2003
			CZ 20022850 A3	12-02-2003
			WO 0164637 A1	07-09-2001
			WO 0162726 A2	30-08-2001
			EP 1265862 A2	18-12-2002
			EP 1263727 A1	11-12-2002
			HU 0204526 A2	28-04-2003
			HU 0300196 A2	28-06-2003
			NO 20023995 A	21-10-2002
			NO 20023997 A	22-10-2002
			US 2003120080 A1	26-06-2003
			US 2003040631 A1	27-02-2003